

PAPER • OPEN ACCESS

Electron-impact excitation of the $(4p^5 5s 5p)^4 S_{3/2}$ quasimetastable state in Rb

To cite this article: V Roman and S Demes 2020 *J. Phys.: Conf. Ser.* **1412** 132012

View the [article online](#) for updates and enhancements.

You may also like

- [Spectral properties of Sb IV from MCDHF calculations](#)
P Jönsson, S Verdebout and G Gaigalas
- [Spectrum and energy levels of quadruply-ionized molybdenum \(Mo V\)](#)
Joseph Reader and Ahmad Tauheed
- [Atomic structure, radiative lifetime and oscillator strength calculations in doubly ionized molybdenum \(Mo iii\)](#)
Pascal Quinet



The Electrochemical Society
Advancing solid state & electrochemical science & technology

243rd ECS Meeting with SOFC-XVIII

More than 50 symposia are available!

Present your research and accelerate science

Boston, MA • May 28 – June 2, 2023

[Learn more and submit!](#)

Electron-impact excitation of the $(4p^55s5p)^4S_{3/2}$ quasimetastable state in Rb

V Roman^{1*} and S Demes²

¹Institute of Electron Physics, National Academy of Sciences of Ukraine, Uzhgorod, 88017, Ukraine

²Institute for Nuclear Research, Hungarian Academy of Sciences, Debrecen, 4026, Hungary

Synopsis The ejected-electron excitation function of the $(4p^55s5p)^4S_{3/2}$ quasimetastable state in rubidium was experimentally investigated by electron-impact in the energy range from the excitation threshold at 16.64 eV to 100 eV. The obtained data at 0.12 eV energy resolution indicate the presence of negative-ion resonances just above excitation threshold. A comparison is made with the optical excitation function of the $(4p^55s5p)^4S_{3/2} \rightarrow (4p^65p)^2P_{3/2}$ transition ($\lambda = 82.37$ nm), measured earlier with a lower energy resolution of ≈ 1.5 eV.

It is known that in alkali atoms (K, Rb, Cs) there is a subclass of p^6 -core excited quartet states which is due to the mixing with doublet states of the same total electronic angular momentum J possess both radiative and electronic decay channels [1]. As a rule, the radiative decay dominates for such quasimetastable states, while autoionization probability remains small. The latter makes difficult the experimental observation of quasimetastable states in ejected-electron spectra [2]. The optical excitation function for the $(4p^55s5p)^4S_{3/2} \rightarrow (4p^65p)^2P_{3/2}$ transition ($\lambda = 82.37$ nm) in Rb was measured earlier with an incident-electron energy resolution of ≈ 1.5 eV [3]. In the present work we obtained the ejected-electron excitation function of the $(4p^55s5p)^4S_{3/2}$ state with an improved 0.12 eV energy resolution by using energy-selective electron beam and by accurate measurement of the ejected-electron spectra in a broad impact energy range.

The apparatus and the experimental procedure were described in details earlier [4]. The ejected-electron spectra were measured at 54.7° observation angle. Due to the overlapping of the $^4S_{3/2}$ line with the high intensity neighboring line $(4p^54d5s)^4P_{1/2}$ with 16.69 eV excitation threshold, the total relative uncertainty is approximately 40% for the most of our data.

Figure 1 shows the measured ejected-electron excitation function of the $(4p^55s5p)^4S_{3/2}$ state and the optical excitation function for $\lambda = 82.37$ nm [3] in an impact-energy range from the 16.64 eV excitation threshold up to 100 eV. Our comparison shows that there is a fairly good agreement between the two types of data. Due to the better energy resolution of the present measurements, the “fine” structure *a-b* is clearly seen in the ejected-electron function just above the excitation threshold. The broad




Figure 1. The electron-impact excitation functions of the $(4p^55s5p)^4S_{3/2}$ state in rubidium: ● - present data; ○ - optical data [3]. The inset shows the threshold region of the present data. The data [3] and our data are normalized with incident-electron energy of 65 eV.

maximum *c* in our data [3] corresponds to the spin-exchange character of the electronic excitation of the $^4S_{3/2}$ state. The notable scatter of experimental data, which is observed between 20 and 25 eV in the ejected-electron excitation function could reflect the combined effect of several overlapping resonances.

In order to obtain the absolute values of the excitation cross sections for the $(4p^55s5p)^4S_{3/2}$ and other p^6 -core excited states in rubidium, we have performed relativistic distorted-wave calculations.

References

- [1] Mendelson A *et al* 1987 *Phys. Rev. A.* **35** 2095
- [2] Borovik A *et al* 2006 *Phys. Rev. A.* **73** 062701
- [3] Bogachev G 1974 *Ukr. Phys. J.* **12** 2061
- [4] Borovik A *et al* 2005 *J. Phys. B* **38** 1081

* E-mail: viktoriyaroman11@gmail.com